

Interim Report
for the period
June 1988 to
January 1989

Thermonuclear Reaction Bibliography with Cross Section Data for Four Advanced Reactions

March 1989

Author:
L. T. Cox, Jr.

AD-A206 915

Approved for Public Release

Distribution is unlimited. The AFAL Technical Services Office has reviewed this report, and it is releasable to the National Technical Information Service, where it will be available to the general public, including foreign nationals.

Air Force
Astronautics
Laboratory

Air Force Space Technology Center
Space Division, Air Force Systems Command
Edwards Air Force Base,
California 93523-5000

DTIC
ELECTE
S 7 APR 1989 D
E


NOTICE

When U.S. Government drawings, specifications, or other data are used for any purpose other than a definitely related Government procurement operation, the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication or otherwise, or in any way licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may be related thereto.

FOREWORD


This interim report was submitted by Larry T. Cox, Jr. documenting work he performed while working at the Air Force Astronautics Laboratory (AFAL), Edwards AFB, CA. AFAL Project Manager is Dr Frank Mead.

This report has been reviewed and is approved for release and distribution in accordance with the distribution statement on the cover and on the DD Form 1473.


FRANKLIN B. MEAD, JR.
Project Manager


CLARENCE J.C. COLEMAN, JR., CAPT, USAF
Chief, Advanced Concepts Branch

FOR THE COMMANDER




REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for Public Release; Distribution is Unlimited		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) AFAL-TR-89-006			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Air Force Astronautics Laboratory		6b. OFFICE SYMBOL (If applicable) LSVF	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Edwards AFB CA 93523-5000			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
PROGRAM ELEMENT NO. 62302F		PROJECT NO. 3058	TASK NO. 00	WORK UNIT ACCESSION NO. J5	
11. TITLE (Include Security Classification) Thermonuclear Reaction Bibliography with Cross Section Data for Four Advanced Reactions (U)					
12. PERSONAL AUTHOR(S) Cox, Larry T. Jr.					
13a. TYPE OF REPORT Interim		13b. TIME COVERED FROM 88.06 TO 89.01		14. DATE OF REPORT (Year, Month, Day) 89,03	
15. PAGE COUNT 35					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
21	06		Thermonuclear reactions		
18	09		Cross Section Data.		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>The Air Force Astronautics Laboratory (AFAL) is exploring the feasibility of fusion-powered propulsion, and is playing a vital role in the push for its development. This report builds on the earlier work of Jim Pass who assembled a bibliography of all known thermonuclear reactions involving reactant particles up to and including B¹¹ (Boron-11). However, in the report presented here, the neutron-induced reactions have been eliminated. A few reactions which are new to the earlier report have been found and are included.</p> <p>This report will form the basis for future reports generated from this study. In these future reports, fuel cycles, reaction rates, nuclear scattering, and ignition temperatures will be addressed. Included in this interim report are cross section data for four of the more attractive, in terms of safety, reactions. Graphs showing the data are also presented herein, including one which compares the cross section data curves of the four reactions. All of the data sets are analyzed and recommendations for future cross section work are made.</p>					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED / UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Franklin B. Mead, Jr.			22b. TELEPHONE (Include Area Code) (805) 275-5540		22c. OFFICE SYMBOL AFAL/LSVF

TABLE OF CONTENTS

SECTION	PAGE
INTRODUCTION, SUMMARY, AND STUDY OVERVIEW	1
THERMONUCLEAR REACTION BIBLIOGRAPHY	3
BACKGROUND	3
LIST OF SYMBOLS	4
PROTON-INDUCED REACTIONS	5
DEUTERON-INDUCED REACTIONS	8
TRITON-INDUCED REACTIONS	11
HELIUM-3-INDUCED REACTIONS	14
HELIUM-4 (ALPHA PARTICLE)-INDUCED REACTIONS	16
LITHIUM-6-INDUCED REACTIONS	17
CROSS SECTIONS FOR PROMISING PROPULSION REACTIONS	18
PROMISING REACTIONS/CRITERIA	18
CROSS SECTION NOTES AND GRAPHS	18
CONCLUSIONS, RECOMMENDATIONS, AND STUDY OUTLOOK	26
REFERENCES	28



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

LIST OF FIGURES

<u>FIGURE</u>	<u>CAPTION</u>	<u>PAGE</u>
1	$\text{He}^3(d,p)\text{He}^4$ Cross Section.....	19
2	$\text{He}^3(\text{He}^3,2p)\text{He}^4$ Cross Section.....	21
3	$\text{Li}^6(p,\text{He}^3)\text{He}^4$ Cross Section.....	22
4	$\text{B}^{11}(p,3\text{He}^4)$ Cross Section.....	23
5	$\text{B}^{11}(p,n)\text{C}^{11}$ Cross Section.....	24
6	Comparative Cross Sections.....	25

INTRODUCTION AND SUMMARY

The Air Force Astronautics Laboratory (AFAL) is exploring the feasibility of fusion-powered propulsion and is continuing to play a vital role in the push for its development. A 3-year study entitled "Fusion Fuel Cross Section" is now underway at the AFAL. For this study, cross section data for advanced fuels is being collected and a computer program will be developed to utilize the information in an extension of the results obtained in the "Barn Book" [Ref. 1]. During the study there will be interim reports published to show progress and to provide up-to-date information with a final report published at the end of the study.

This interim report builds on the earlier work of Jim Pass [Ref. 2] who assembled a bibliography of all known thermonuclear reactions involving reactant particles up to and including B^{11} (Boron-11). In this report the neutron-induced reactions have been eliminated. In later work, reaction specifics including cross sections, reaction parameters, and curve-fit equations will be compiled, developed, and tabulated for the reactions. Throughout the study conclusions will be made concerning which data are most accurate and which reactions seem to be the best candidates for propulsion fuels, thus giving researchers an idea of where to focus their work. The reports which evolve will serve as valuable references for anyone studying advanced thermonuclear reactions and their respective characteristics.

Many reasons justify Air Force interest in advanced fusion propulsion. In particular, the limits of chemical systems are driving factors to search for a new alternative. Several aspects of fusion push for its development. These include high specific impulse (1500 to 10,000 seconds), limited neutron and gamma-ray production, high thrust, and environmentally safe exhaust products. Such possibilities will enable space missions never before realized and will greatly enhance or make more efficient current Air Force missions.

This report includes reaction lists which have been cross-referenced extensively to insure accuracy. Cross section data have been compiled to produce plots for some of the reactions viewed as promising because of negligible production of harmful radiation. An assessment of the data which is currently available for each of these chosen reactions is included. Additional data will be continually collected for all the reactions listed. Ultimately the final report will include comprehensive information for each reaction.

Compiling the reaction list involved a re-examination of most of the references used in the earlier report [Ref. 2] as well as a detailed inspection of additional thermonuclear reaction references. If a reaction listed in the earlier report could not be found in available resources, the source referenced in the earlier report is referenced once again. As additional sources become available in the future, the reactions in question will be checked for their accuracy. The search for, and interpretation of, cross section data involved contacting fusion experts at various universities and national laboratories. Two national laboratories were of particular help in this effort. Brookhaven National Laboratory [Ref. 3] provided access to its huge nuclear cross section data file, and Los Alamos National Laboratory [Ref. 4] contributed its latest cross section data, which was fitted using R-Matrix analysis. In addition, the Fusion Studies Laboratory of the University of Illinois at Urbana-Champaign was timely in sending up-to-date data for the $\text{He}^3\text{-He}^3$ reaction.

Some interesting developments have emerged. Differences between previously published data and the data available here are presented, and other notable points are discussed in the section entitled "CROSS SECTION NOTES AND GRAPHS." An assessment of the available data and recommendations for further research accompany it. Future aims of the study are roughly outlined as well.

THERMONUCLEAR REACTION BIBLIOGRAPHY

On the next few pages are lists of possible fuels for fusion propulsion systems. Neutron-induced reactions have been excluded from the bibliography, as the aim of this interim report is to look closely at cross section data for a few hopeful reactions. Later in this study, a report which includes the neutron-based reactions may be published as a tool for determining the constituent reactions of a fuel cycle. Until such a time, consult the earlier bibliography [Ref. 2] for this information. Neutron reactions are also documented in other sources, such as S. Flugge's work [Ref. 5].

BACKGROUND

Since the 1950's, no large efforts have been made to discover thermonuclear reactions in the isotopic range between H^1 and B^{11} . Thus, the works of S. Flugge [Ref. 5] and others remain important sources to this day. Current efforts exploit these known reactions, but do not attempt to find new ones. Thus, the compilation presented here should not be viewed as complete. There may be additional reactions remaining to be found, and increased interest in fusion may lead to their discovery. It should be noted that many independent data sources comprise Ref. 3, a fact which should explain seemingly conflicting information in the following pages. This may also surface in reference to other sources.

LIST OF SYMBOLS [Ref. 2]

Q = $Q(\text{MeV})$ values taken from Ref. 2 or calculated using the mass defect formula:

$$Q = [(m_a + m_b) - (m_c + m_d + \dots)] \cdot C$$

$$C = 931.481 \text{ MeV/amu}$$

m_a = mass of target particle

m_b = mass of incident particle

m_c, m_d, \dots = masses of reaction products

The two notations for a thermonuclear reaction are related as follows:

$$a + b \rightarrow c + d + \dots = a(b,c)d + \dots,$$

where the species within the parentheses are the particles of smallest reactant mass (b) and smallest initial product mass (c). The second notation is used in conjunction with the cross section graphs.

Other symbols used:

$*$ = isotope in excited state

γ = gamma-ray

e^- = electron

e^+ = positron

n = neutron

p = proton

d = deuteron

t = triton

\rightarrow = this arrow separates reactants and products

\rightarrow = reaction product preceding this arrow decays as follows

$\text{He}^{3,4,5,6}$ = Helium-3,4,5,6

$\text{Li}^{5,6,7,8}$ = Lithium-5,6,7,8

$\text{Be}^{7,8,9,10}$ = Beryllium-7,8,9,10


$\text{B}^{9,10,11,12}$ = Boron-9,10,11,12

$\text{C}^{10,11,12,13,14}$ = Carbon-10,11,12,13,14

$\text{N}^{13,14}$ = Nitrogen-13,14

NOTE: All reactions are listed by increasing atomic number of the target nuclide. For a reaction with more than one branch, reactant particles are listed only once, with each branch's products given on a separate line.











PROTON-INDUCED REACTIONS



$p + \text{Be}^7 \rightarrow g + \text{B}^8 + 0.138 \text{ MeV [Ref. 17]}$
 $p + \text{Be}^9 \rightarrow n + p + \text{Be}^8 \rightarrow 2\text{He}^4 - 1.67 \text{ MeV [Ref. 5]}$
 $\rightarrow n + \text{B}^9 \rightarrow p + 2\text{He}^4 - 1.852 \text{ MeV [Ref. 5]}$
 $\rightarrow \text{He}^4 + \text{Li}^6 + 2.1 \text{ MeV [Refs. 19,20]}$
 $\quad + 2.125 \text{ MeV [Ref. 9]}$
 $\quad + 2.126 \text{ MeV [Ref. 5]}$
 $\quad + 2.13 \text{ MeV [Ref. 14]}$
 $\text{He}^4(1.275) + \text{Li}^6(0.850) = 2.125 \text{ MeV [Ref. 18]}$
 $\rightarrow g + \text{B}^{10} \text{ [Ref. 5]}$
 $\quad + 6.585 \text{ MeV [Ref. 17]}$
 $\rightarrow d + 2\text{He}^4 + 0.56 \text{ MeV [Ref. 14]}$
 $\quad + 0.7 \text{ MeV [Refs. 19,20]}$
 $\rightarrow d + \text{Be}^8 \rightarrow 2\text{He}^4 \text{ [Ref. 5]}$
 $\quad + 0.651 \text{ [Ref. 17]}$
 $\rightarrow t + \text{Be}^7 - 12.17 \text{ MeV [Ref. 5]}$
 $\rightarrow \text{B}^{10}(6.88) - .295 \text{ MeV [Ref. 20]}$
 $p + \text{B}^{10} \rightarrow \text{He}^4 + \text{Be}^7 + 1.148 \text{ MeV [Ref. 5]}$
 $\quad + 1.147 \text{ MeV [Ref. 9]}$
 $\rightarrow n + \text{C}^{10} \rightarrow \text{B}^{10} + e^+ - 4.35 \text{ MeV [Ref. 5]}$
 $\rightarrow g + \text{C}^{11} \rightarrow \text{B}^{11} + e^+ \text{ [Ref. 5]}$
 $\quad + 8.690 \text{ MeV [Ref. 17]}$
 $\rightarrow d + \text{B}^9 \rightarrow p + 2\text{He}^4 - 5.877 \text{ MeV [Ref. 5]}$
 $\rightarrow t + \text{B}^8 \rightarrow e^+ + \text{Be}^8 \rightarrow 2\text{He}^4 - 1.00 \text{ MeV [Ref. 5]}$
 $\rightarrow \text{He}^3 + \text{Be}^8 \rightarrow 2\text{He}^4 \text{ [Ref. 5]}$
 $\quad - 0.442 \text{ MeV [Ref. 17]}$

$p + B^{11} \rightarrow He^4 + Be^8(\text{ground state}) \rightarrow 2He^4$ [Ref. 21]
 $\rightarrow He^4 + Be^{8*}$ [Ref. 21]
 $\rightarrow He^4 + Be^8 \rightarrow 2He^4 + 8.585 \text{ MeV}$ [Ref. 5]
 $\rightarrow 3He^4$ [Ref. 21]
 $\quad + 8.7 \text{ MeV}$ [Refs. 15,19,21,22,23,24,25]
 $\quad + 8.68 \text{ MeV}$ [Ref. 14]
 $3He^4(2.888 \text{ each}) = 8.664 \text{ MeV}$ [Ref. 9]
 $\rightarrow n + C^{11} \rightarrow B^{11} + e^+ - 2.762 \text{ MeV}$ [Ref. 5]
 $\rightarrow g + C^{12}$ [Ref. 5]
 $\quad + 15.956 \text{ MeV}$ [Ref. 17]
 $\rightarrow d + B^{10} - 9.10 \text{ MeV}$ [Ref. 5]
 $\rightarrow C^{12*} + 16.11 \text{ MeV}$ [Ref. 26]
 $\quad + 16.58 \text{ MeV}$ [Ref. 26]














DEUTERON-INDUCED REACTIONS



$d + \text{He}^3 \rightarrow p^*(14.68) + \text{He}^4(3.67) = 18.35 \text{ MeV}$ [Ref. 7]
 $d + \text{He}^4 \rightarrow n + p + \text{He}^4 - 2.2 \text{ MeV}$ [Ref. 6]
 $\rightarrow p + \text{He}^5 \rightarrow n + \text{He}^4 - 2.22? \text{ MeV}$ [Ref. 3]
 $d + \text{Li}^6 \rightarrow g + n + \text{Be}^7 + 3.38? \text{ MeV}$ [Ref. 3]
 $\rightarrow n + \text{He}^3 + \text{He}^4 + 1.72 \text{ MeV}$ [Ref. 6]
 $\quad + 1.796 \text{ MeV}$ [Refs. 9,10]
 $\quad + 1.8 \text{ MeV}$ [Refs. 11,12,19]
 $n(1.134) + \text{He}^3(0.378) + \text{He}^4(0.284) = 1.796 \text{ MeV}$ [Refs. 2,8]
 $\rightarrow n + \text{Be}^7 + 3.34 \text{ MeV}$ [Ref. 6]
 $\quad + 3.40 \text{ MeV}$ [Refs. 5,11,12,16,19]
 $n(2.957) + \text{Be}^7(0.423) = 3.380 \text{ MeV}$ [Refs. 8,9]
 $\rightarrow n + \text{Be}^7 \rightarrow n + \text{He}^3 + \text{He}^4 + 6.28? \text{ MeV}$ [Ref. 13]
 $\rightarrow p + \text{Li}^7 + 5.0 \text{ MeV}$ [Refs. 11,12,16,19]
 $\quad + 5.02 \text{ MeV}$ [Ref. 6]
 $\quad + 5.026 \text{ MeV}$ [Refs. 9,10]
 $\quad + 5.027 \text{ MeV}$ [Ref. 5]
 $p(4.398) + \text{Li}^7(0.628) = 5.026 \text{ MeV}$ [Refs. 8,18]
 $\rightarrow p + \text{Li}^7 \rightarrow p + t + \text{He}^4 + 4.73? \text{ MeV}$ [Ref. 13]
 $\rightarrow p + \text{Li}^7 \rightarrow g + \text{Li}^7 + (4.54 + 0.45) \text{ MeV}$ [Ref. 6]
 $\rightarrow p + t + \text{He}^4 + 2.557 \text{ MeV}$ [Ref. 17]
 $\quad + 2.561 \text{ MeV}$ [Refs. 9,10]
 $\quad + 2.6 \text{ MeV}$ [Refs. 11,12,16,19]
 $\rightarrow t + \text{Li}^5 \rightarrow p + \text{He}^4 + (0.9 + 1.6) \text{ MeV}$ [Ref. 6]
 $t(0.539) + p(1.618) + \text{He}^4(0.404) = 2.561 \text{ MeV}$ [Refs. 8,18]
 $\rightarrow \text{He}^3 + \text{He}^5 \rightarrow n + \text{He}^4 + (0.8 + 1.0) \text{ MeV}$ [Ref. 16]
 $\rightarrow 2\text{He}^4 + 22.373 \text{ MeV}$ [Ref. 10]
 $\quad + 22.39 \text{ MeV}$ [Ref. 5]
 $\quad + 22.4 \text{ MeV}$ [Refs. 6,11,12,15,16,19,22]
 $2\text{He}^4(11.187 \text{ each}) = 22.374 \text{ MeV}$ [Refs. 8,9,18]
 $d + \text{Li}^7 \rightarrow g + 2n + \text{Be}^7 - 3.87? \text{ MeV}$ [Ref. 3]
 $\rightarrow g + p + \text{Li}^8 \rightarrow e^- + \text{Be}^8 \rightarrow 2\text{He}^4 + 15.4? \text{ MeV}$ [Ref. 3]
 $\rightarrow 2\text{He}^4 + 15.9? \text{ MeV}$ [Ref. 3]
 $\rightarrow 2n + \text{Be}^7 - 3.869 \text{ MeV}$ [Ref. 17]

$d + Li^7 \rightarrow n + Be^8 \rightarrow 2He^4 + 15.0 \text{ MeV [Refs. 5,6]}$
 $n(10.082) + 2He^4(2.521 \text{ each}) = 15.124 \text{ MeV [Ref. 18]}$
 $\rightarrow p + Li^8 \rightarrow e^- + Be^8 \rightarrow 2He^4 + (-0.26 + 16.0) \text{ MeV [Ref. 6]}$
 $\rightarrow t + Li^6 = 0.995 \text{ MeV [Ref. 6]}$
 $\rightarrow He^4 + He^5 \rightarrow n + He^4 + 14.2 \text{ MeV [Ref. 5]}$
 $\rightarrow Be^9 + 16.674 \text{ MeV [Ref. 26]}$
 $d + Be^7 \rightarrow p + 2He^4 + 16.7 \text{ MeV [Ref. 16]}$
 $+ 16.766 \text{ MeV [Ref. 17]}$
 $p(11.179) + 2He^4(2.795 \text{ each}) = 16.769 \text{ MeV [Ref. 18]}$
 $d + Be^9 \rightarrow \alpha + B^{11} + 15.87 \text{ MeV [Ref. 3]}$
 $\rightarrow p + Be^{10} + 4.588 \text{ MeV [Ref. 5]}$
 $\rightarrow n + B^{10} + 4.35 \text{ MeV [Ref. 5]}$
 $\rightarrow t + Be^8 \rightarrow 2He^4 + 4.687 \text{ MeV [Ref. 3]}$
 $\rightarrow He^4 + Li^7 + 7.153 \text{ MeV [Ref. 5]}$
 $d + B^{10} \rightarrow p + B^{11} + 9.235 \text{ MeV [Ref. 5]}$
 $\rightarrow n + C^{11} \rightarrow B^{11} + e^+ + 6.6 \text{ MeV [Ref. 5]}$
 $\rightarrow He^4 + Be^8 \rightarrow 2He^4 + 17.86 \text{ MeV [Ref. 5]}$
 $d + B^{11} \rightarrow n + C^{12} + 13.8 \text{ MeV [Ref. 5]}$
 $\rightarrow 2n + C^{11} = 4.997 \text{ MeV [Ref. 3]}$
 $\rightarrow p + B^{12} \rightarrow e^- + C^{12} + 1.137 \text{ MeV [Ref. 5]}$
 $\rightarrow He^4 + Li^8 \rightarrow e^- + Be^9 \rightarrow 2He^4$
 $+ 1.137 \text{ MeV [Ref. 5]}$
 $\rightarrow He^4 + Be^9 + 8.018 \text{ MeV [Ref. 5]}$

TRITON-INDUCED REACTIONS



$t + Li^6 \rightarrow g + p + Li^8 \rightarrow e + Be^8 \rightarrow 2He^4 + 16.47 \text{ MeV [Ref. 3]}$
 $\rightarrow 2He^4 + 16.97 \text{ MeV [Ref. 3]}$
 $\rightarrow d + Li^7 + 0.994 \text{ MeV [Ref. 17]}$
 $\quad + 0.995 \text{ MeV [Ref. 6]}$
 $\quad + 1.0 \text{ MeV [Refs. 12, 16]}$
 $\rightarrow d + Li^{7*} \rightarrow g + Li^7 + (0.509 + 0.45) \text{ MeV [Ref. 6]}$
 $\rightarrow 2n + Be^7 + 2.876 \text{ MeV [Ref. 17]}$
 $\rightarrow p + Li^8 + 0.8 \text{ MeV [Ref. 16]}$
 $\rightarrow e + 2He^4 + 0.800 \text{ MeV [Ref. 6]}$
 $\rightarrow n + Be^8 \rightarrow 2He^4 + 16.0 \text{ MeV [Ref. 6]}$
 $\rightarrow n + 2He^4 + 16.0 \text{ MeV [Ref. 6]}$
 $\quad + 16.1 \text{ MeV [Ref. 16]}$
 $\rightarrow g + Be^9 + 17.77 \text{ MeV [Ref. 5]}$
 $\rightarrow He^4 + He^5 \rightarrow n + He^4 + 16.17 \text{ MeV [Ref. 5]}$
 $t + Li^7 \rightarrow g + d + Li^8 \rightarrow e + Be^8 \rightarrow 2He^4 + 11.47 \text{ MeV [Ref. 3]}$
 $\rightarrow 2He^4 + 11.97 \text{ MeV [Ref. 3]}$
 $\rightarrow 2n + 2He^4 \text{ [Ref. 6]}$
 $\rightarrow 2n + 2He^4 + 8.88 \text{ MeV [Ref. 6]}$
 $\quad 2n(6.049 \text{ each}) + 2He^4(1.512 \text{ each}) = 15.122 \text{ MeV [Ref. 18]}$
 $\rightarrow n + Be^9 + 10.4 \text{ MeV [Ref. 12]}$
 $\quad + 10.52 \text{ MeV [Ref. 6]}$
 $\rightarrow 2n + Be^8 \rightarrow 2He^4 + 8.83 \text{ MeV [Ref. 6]}$
 $\rightarrow 2n + 2He^4 + 8.8 \text{ MeV [Ref. 12]}$
 $\rightarrow n + He^4 + He^5 \rightarrow n + He^4 + (8.08 + 1.0) \text{ MeV [Ref. 6]}$
 $\rightarrow p + Li^9 \rightarrow e + Be^9 + 10.77 \text{ MeV [Ref. 3]}$
 $\rightarrow n + 2He^4 + 9.657 \text{ MeV [Ref. 3]}$
 $\rightarrow d + Li^8 \rightarrow e + Be^8 \rightarrow 2He^4 + 11.47 \text{ MeV [Ref. 3]}$
 $\rightarrow 2He^4 + 11.97 \text{ MeV [Ref. 3]}$
 $\rightarrow He^4 + He^6 \rightarrow e + Li^6 + 9.83 \text{ MeV [Ref. 6]}$

$$t + \text{Be}^7 \rightarrow p + n + 2\text{He}^4 \text{ [Ref. 2]} \\ p(4.204) + n(4.204) + 2\text{He}^4(1.051 \text{ each}) = 10.510 \text{ MeV [Ref. 2]}$$

$$t + \text{Be}^9 \rightarrow n + \text{B}^{11} + 9.56? \text{ MeV [Ref. 3]}$$

$$t + \text{B}^{10} \rightarrow \text{Be}^7 + \text{He}^6 \rightarrow e^- + \text{Li}^6 - 3.36? \text{ MeV [Ref. 3]}$$

$$\rightarrow p + \text{B}^{12} \rightarrow e^- + \text{C}^{12} + 19.2? \text{ MeV [Ref. 3]}$$

$$\rightarrow \text{He}^4 + \text{Li}^8 \rightarrow e^- + \text{Be}^8 \rightarrow 2\text{He}^4 + \\ 11.9? \text{ MeV [Ref. 3]}$$











$$\rightarrow 2\text{He}^4 + 12.4? \text{ MeV [Ref. 3]}$$

$$\rightarrow p + \text{He}^4 + \text{Li}^8 \rightarrow e^- + \text{Be}^8 \rightarrow 2\text{He}^4 + 11.9? \text{ MeV [Ref. 3]}$$

$$\rightarrow 2\text{He}^4 + 12.4? \text{ MeV [Ref. 3]}$$

$$t + \text{B}^{11} \rightarrow 2\text{He}^4 + \text{He}^6 \rightarrow e^- + \text{Li}^6 - 39.5? \text{ MeV [Ref. 3]}$$

HELIUM-3-INDUCED REACTIONS




$\text{He}^3 + \text{Li}^7 \rightarrow n + \text{B}^9 \rightarrow p + \text{Be}^8 \rightarrow 2\text{He}^4 + (9.3 + 0.3) \text{ MeV [Ref. 6]}$
 $\rightarrow n + p + \text{Be}^8 \rightarrow 2\text{He}^4 + (9.5 + 0.1) \text{ MeV [Ref. 6]}$
 $n(3.852) + p(3.852) + 2\text{He}^4(1.512) = 10.728 \text{ MeV [Ref. 18]}$
 $\rightarrow p + \text{Be}^9 + 11.2 \text{ MeV [Ref. 6]}$
 $\rightarrow d + \text{Be}^8 \rightarrow 2\text{He}^4 + (11.7 + 0.1) \text{ MeV [Ref. 6]}$
 $\rightarrow t + \text{Be}^7 - 0.881? \text{ MeV [Ref. 3]}$
 $\rightarrow 2p + 2\text{He}^4 \text{ [Ref. 18]}$
 $2p(4.510 \text{ each}) + 2\text{He}^4(1.127 \text{ each}) = 11.274 \text{ MeV [Ref. 18]}$
 $\rightarrow \text{He}^4 + \text{Li}^6 + 13.3 \text{ MeV? [Ref. 5]}$




















$\text{He}^3 + \text{Be}^9 \rightarrow n + \text{He}^4 + \text{Be}^7 + 0.014? \text{ MeV [Refs. 3, 26]}$
 $\rightarrow 2n + \text{C}^{10} \rightarrow e^+ + \text{B}^{10} - 2.43? \text{ MeV [Ref. 3]}$
 $\rightarrow p + \text{B}^{11} + 10.3? \text{ MeV [Ref. 5]}$
 $\rightarrow p + \text{B}^{11*} + 10.32 \text{ MeV [Ref. 26]}$
 $\rightarrow n + \text{C}^{11} \rightarrow e^+ + \text{B}^{11} + 9.02? \text{ MeV [Ref. 5]}$
 $\rightarrow \text{He}^4 + \text{Be}^8 \rightarrow 2\text{He}^4 + 19.0? \text{ MeV [Ref. 5]}$
 $\rightarrow \text{He}^4 + \text{Be}^{8*} + 18.94 \text{ MeV [Ref. 26]}$

$\text{He}^3 + \text{B}^{10} \rightarrow p + \text{C}^{12} + 19.7? \text{ MeV [Ref. 5]}$
 $\rightarrow p + \text{C}^{12*} + 19.67 \text{ [Ref. 26]}$
 $\rightarrow t + \text{C}^{10} \rightarrow e^+ + \text{B}^{10} - 0.530? \text{ MeV [Ref. 3]}$
 $\rightarrow \text{He}^4 + \text{B}^9 \rightarrow p + 2\text{He}^4 + 12.4? \text{ MeV [Ref. 5]}$
 $\rightarrow d + \text{C}^{11} \rightarrow e^+ + \text{B}^{11} + 4.73? \text{ MeV [Ref. 5]}$
 $\rightarrow \text{Li}^6 + \text{Be}^7 - 2.87? \text{ MeV [Ref. 3]}$
 $\rightarrow p + \text{C}^{13} + 13.2? \text{ MeV [Ref. 5]}$
 $\rightarrow d + \text{C}^{12} + 10.6? \text{ MeV [Ref. 5]}$

HELIUM-4 (ALPHA-PARTICLE)-INDUCED REACTIONS



LITHIUM-6-INDUCED REACTIONS



CROSS SECTIONS FOR PROMISING PROPULSION REACTIONS

PROMISING REACTIONS/CRITERIA

On the following pages the cross sections for some of the more attractive reactions for propulsion are investigated. Some characteristics of an ideal reaction for a propulsion system include the following:

1. No neutron production.
2. No gamma-ray production.
3. Exothermic in nature.
4. Achievable ignition.

The reactions examined were chosen mainly because they produce minimal amounts of harmful radiation (criteria 1 and 2). All of the reactions are exothermic (3), which is an obvious propulsion system requirement. Achieving ignition (4) is the sticking point which should be resolved as containment technology advances.

Four reactions are considered: 1) $d-He^3$, 2) He^3-He^3 , 3) $p-Li^6$, and 4) $p-B^{11}$. The $d-He^3$ reaction will likely be the first of these to be utilized. It produces some neutrons, but methods to reduce them, such as spin polarization, are being researched. He^3-He^3 would be the ideal choice for propulsion because it is completely aneutronic. At this time, however, it seems to be an unlightable reaction, as illustrated by the compiled data. Both $p-Li^6$ and $p-B^{11}$ come close to being aneutronic, but ignition seems to be the major obstacle because of the high bremsstrahlung and cyclotron radiation energy loss.

CROSS SECTION NOTES AND GRAPHS

Graphed data for the $d-He^3$ reaction are shown in Figure 1. This reaction has been documented in many sources and is the object of a number of research projects. As can be seen, the curve is well-defined by the closely-agreeing sources of data. There is some discrepancy as to where the maximum occurs, as it varies from 720 to 818 mb at the peak energy. Methods now need to be devised to minimize neutron production. It is currently planned for neutron-producing branches of this reaction to be considered in a future report. The remaining concern is being able to achieve the approximately 0.45 MeV temperature at which the reaction reaches its resonance peak. It is at this point where the the highest probability of the maximum reaction rate is obtained. If not obvious from the graph, the dashed line corresponds to the curve obtained from the "Barn Book" [Ref. 1].




Fig. 1. $\text{He}^3(\text{d},\text{p})\text{He}^4$ Cross Section

The cross section data for $\text{He}^3\text{-He}^3$ are shown in Figure 2. Data points found by Good in 1954 are shown as boxes. The curve shown was determined by Dwarakanath's 1971 data (dots) and Pritzker's 1976 data (X's) [Ref. 32]. No curve peaks (maxima) are readily apparent--the only possibility being along the level curve portion in the range of 8 to 10.4 MeV. A temperature of this magnitude is beyond the limits of present confinement technology. Thus, it is apparent that this reaction may not be ignitable, at least not in the near future. Even so, additional data may provide a breakthrough by revealing a peak in the cross section curve along the level portion.

Another reaction which needs more data to adequately define the cross section curve is the p-Li^6 reaction, shown in Figure 3. The latest findings [Ref. 4] show a well-fit curve for the reaction. For the energy range common to the two sets of data, the shape of the curve is similar. Additional data for the peak region is needed so a precise cross section value can be determined for use in experiments. There is also a gamma-ray producing branch, but no significant data was found. The Ref. 3 curve is shown without data points for clarity.

Figures 4 and 5 show the p-B^{11} reaction. The favorable branch, $\text{B}^{11}(\text{p},3\text{He}^4)$, is represented in Figure 4. According to recent data collected at Los Alamos [Ref. 33] and graphs from Ref. 34, a resonance may exist at a lower energy than previously noted in such sources as the Barn Book [Ref. 1]. Though not shown on the Ref. 4 curve, this unproven spike appears at 0.162 MeV. Otherwise, the general shapes of the two curves are roughly similar. At lower energies, the differences begin to grow quite large. In Figure 5 the less favorable $\text{B}^{11}(\text{p},\text{n})\text{C}^{11}$ branch is detailed. The data set denoted by the "X's" is small and departs decidedly from the other data curves. One point (0.2 MeV, 2.5 mb) from this set was omitted on the graph in order to narrow the spectrum of temperatures shown, thereby increasing detail of the curves. The other two data sets appear as though they could blend into a single curve, if the higher-energy set is viewed as a continuance of the first. If Figure 5 is compared to the favorable branch in Figure 4, it can be concluded that the unwanted branch's cross sections fall off rapidly as energy drops, provided the "X" data points are ignored. It would seem that by staying at temperatures below 3 MeV, the neutron-producing branch would not become a significant factor.

Figure 6 is a graph showing the comparative cross sections for each of the four fuels. The $\text{He}^3\text{-He}^3$ curve is a reprint of the earlier graph formed from the Dwarakanath and Pritzker data. All other curves in Figure 6 show the Los Alamos R-matrix data analysis results [Refs. 4,33]. Some data points have been omitted from Figure 6 to maintain clarity; d-He^3 is shown without them altogether. For p-B^{11} the accepted maximum is reached at 0.625 MeV with a cross-section of 800 mb. D-He^3 exhibits a peak of approximately 847 mb at 0.426 MeV. Finally, p-Li^6 peaks at a stable 219 mb from 1.76 to 1.81 MeV. In the future a more precise curve will be determined for each fuel, after additional cross-section data is obtained. Brookhaven National Laboratory [Ref. 3], Los Alamos National Laboratory [Refs. 4,33], the Barn Book [Ref. 1], and the AEP Barnbook [Ref. 32] were the main sources of data for these graphs. It should be noted that data from the Barn Book are shown as curves without points in Figs. 1 and 4, as was done in the reference itself.




Fig. 2. $\text{He}^3(\text{He}^3, 2p)\text{He}^4$ Cross Section




Fig. 3. $\text{Li6}(p,\text{He3})\text{He4}$ Cross Section




Fig. 4. $B_{11}(p,3He_4)$ Cross Section




Fig. 5. $B_{11}(p,n)C_{11}$ Cross Section




Fig. 6. Comparative Cross Sections

CONCLUSIONS, RECOMMENDATIONS, AND STUDY OUTLOOK

During this initial phase of the study, a great deal has been learned about thermonuclear reactions and the meanings of their cross sections. Already several conclusions about available cross section information can be drawn. First, most of the references that have been consulted were generally 10 to 20 years old, some even older. This may be due to the fact that only recently has a genuine effort been started to make fusion space propulsion a reality. However, most of the data presented here is less than 10 years old. Secondly, the safety involved with an advanced fuel that is a neutronic has been, and will continue to be, a major factor in the push for its development. Finally, it is surprising that no significant work has been done since the 1950's to discover new reactions using the simple bombardment technique. It should be fairly obvious, though, that the number of reactions is quite plentiful, and the number which may await discovery should not be large in number.

The conclusions basically center around the cross section information which has been published and is presented here. The $d\text{-He}^3$ initial reaction's cross section data from many sources agree well, so there is a solid foundation for anyone attempting to ignite it. The other reactions which have been included here are not so well defined. $\text{He}^3\text{-He}^3$ needs much more research in order to truly determine its feasibility. Other than unknown resonance peaks, a problem which needs consideration is the unavailability of He^3 for reaction. Recent studies [Refs. 35,36,37] indicate the moon to be a potential source of He^3 . This makes its labeling as a futuristic fuel quite appropriate. $p\text{-Li}^6$ and $p\text{-B}^{11}$ need additional sources of cross section data to determine if resonance peaks exist at energies below those currently indicated by cross section data. Each of them has a harmful radiation-producing branch that must be investigated in more detail. This will allow researchers to be sure that the cross sections of these unwanted reactions are negligible when compared with the cross sections of the more favorable branches at ignition energies.

From these conclusions some recommendations can be presented for additional research in the cross section field. Spin-polarization and other methods to reduce neutron production in the $d\text{-He}^3$ reaction should continue to be addressed, as this reaction appears to be an excellent candidate due to the closely-agreeing data and low ignition energy it possesses. Mining the moon and the outer planets for reactant fuels such as He^3 should continue to be investigated for feasibility. If the He^3 reactions are found to be ignitable, a plan for obtaining the He^3 needed would already have a great start. Precise measurements for each reaction near its lowest-energy resonance peak should be one focus of additional research in the area. Another focal point should be the cross-section measurements for the radioactive particle-producing branches of each reaction. In the case of $p\text{-B}^{11}$, the possible lower energy resonance needs to be investigated more fully, as it has encountered mixed reviews concerning its validity. As is obvious, a lower resonance energy will enable fusion to become a reality sooner than expected and will relieve some of the stress on containment technology.

In the future, this study will involve obtaining cross sections and reactivities for additional advanced reactions. All of this data will be fitted to curves and presented along with discussion about which sets of data are best. Based on this data, and with parameters such as nuclear scattering and various sources of radiation loss taken into account, new ignition temperatures will be calculated for each of the reactions. Additional sources for reactions will continue to be consulted, so an up-to-date reaction bibliography will exist. Eventually, complete fuel cycles will be detailed to provide complete information for researchers attempting to develop reactions into feasible fuel systems. Much of the new information will be the basis for computer programs written to aid in the calculation of new ignition temperatures and reactivities.

REFERENCES

- 1) Miley, G. H., Towner, M., and Ivich, M., Fusion Cross Sections and Reactivities (the "Barn Book"), Report C00-2218-17, University of Illinois, C-U Campus, Nuclear Engineering Program, Urbana IL, 17 June 1974.
- 2) Pass, J. J., Jr., Fusion and Advanced Fuel Reaction Bibliography-- Particle Reactions from H^I to B^{II}, AFAL-TR-87-073, Air Force Astronautics Laboratory, Edwards Air Force Base CA, August 1987.
- 3) Cross Section Data for Thermonuclear Reactions Involving Particles in the Atomic Range Z = 1 to 105, EXFOR Format, National Nuclear Data Center, Brookhaven National Laboratory, Upton, Long Island NY, 1987.
- 4) Hale, G. M., Fusion Cross Sections in ENDF-like Format, University of California, Los Alamos National Laboratory, Los Alamos NM, 5 March 1981.
- 5) Flugge, S., Encyclopedia of Physics--Nuclear Reactions 1, Vol. 40, Springer-Verlag oHG, Berlin-Göttingen-Heidelberg, pp. 63-151, 1957.
- 6) Crocker, V. S., Blow, S., and Watson, C. J. H., Nuclear Cross Section Requirements for Fusion Reactors, CLM-P240, Culham Laboratory, Abingdon, Berkshire, England, 1970.
- 7) Ashworth, C. P., A Utility View of Fusion, EPRI ER-536-SR, Proceedings of the Review Meeting on Advanced-Fuel Fusion, edited by C. K. Choi, Electrical Power Research Institute, Palo Alto CA, pp. 23-37, June 1977.
- 8) Sharp, R. D., and McNally, J. R., Jr., DI and Advanced Fusion Fuels Computer Code, ORNL-TM-50, National Technical Information Service, Springfield VA, p. 25, 9 December 1975.
- 9) McNally, J. R., Jr., "Physics of Fusion Fuel Cycles," Nuclear Technology/Fusion, Vol. 2, pp. 2-28, January 1982.
- 10) McNally, J. R., Jr., Reactivity of Advanced Fusion Fuels, ORNL-TM-4647, Oak Ridge National Laboratory, Oak Ridge TN, 1974.
- 11) McNally, J. R., Jr., Advanced Fuels for Nuclear Fusion Reactors, Conf-750303-26, Oak Ridge National Laboratory, Oak Ridge TN, 7 May 1975.
- 12) McNally, J. R., Jr., Nuclear Fusion Chain Reaction Applications in Physics and Astrophysics, IAEA/SM-170/49, International Atomic Energy Agency, Vienna, Austria, pp. 4-12, 1973.

- 13) Holland, R. E., et al., Nuclear Cross Sections for Light Ions on Li^6 , IEEE Transactions on Nuclear Science, Vol. NS-28, No. 2, Physics Division, Argonne National Laboratory, Argonne IL, April 1981.
- 14) Hirshfield, J. L., Fusion Reactors Based on $\text{Li}^6(\text{p}, \text{He}^4)\text{He}^3$, EPRI-536-SR, Proceedings of the Review Meeting on Advanced-Fuel Fusion, edited by C. K. Choi, Electric Power Research Institute, Palo Alto CA, pp. 262-4, June 1977.
- 15) Westinghouse Fusion Power Systems--1978 Calendar and Reference Book, Westinghouse Electric Corp., Pittsburgh PA, 1978.
- 16) McNally, J. R., Jr., "Fusion Chain Reactions--I," Nuclear Fusion, Vol. 11, pp. 187-93, 1971.
- 17) Feldbacher, R., Heindler, M., and Miley, G., Requirements for Charged Particle Light Isotopes Reaction Data for Advanced Fuel Cycles, Including Two-Step Reaction Mechanism, Alternate Energy Physics Program, Technical University Graz, Austria, and University of Illinois, 1986.
- 18) McNally, J. R., Jr., Fusion Reactivity Graphs and Tables for Charged Particle Reactions, ORNL-TM-6914, Oak Ridge National Laboratory, Oak Ridge TN, August 1979.
- 19) Miley, G. H., Fusion Energy Conversion, American Nuclear Society, ANS Publications, Hinsdale IL, 1976.
- 20) McNally, J. R., Jr., "Autocatalytic Burning of Li^6/D Nuclear Fuel via the $\text{p} + \text{Be}^9$ Resonance Reaction," Nuclear Fusion, Vol. 11, pp. 554-6, 1971.
- 21) "Advanced Fusion Power: A Preliminary Assessment," National Academy Press, Washington, D. C., pp. 3-5, 1987.
- 22) Book, D. L., NRL Plasma Formulary, NRL Publication 0084-4040.
- 23) Dingle, D. A., "Fusion Power," Chemical and Engineering News, 2 April 1979.
- 24) Miley, G. H., Comments about p-B¹¹ Ignition, EPRI-536-SR, Proceedings of the Review Meeting on Advanced-Fuel Fusion, edited by C. K. Choi, Electric Power Research Institute, Palo Alto CA, June 1977.
- 25) Committee on Aneutronic Fusion Power, Phase 1, Air Force Studies Board, Commission on Engineering and Technical Systems, National Research Council, National Academy of Sciences, 5 March 1987.
- 26) McNally, J. R., Jr., Nuclear Fusion Resonance Reactions of Possible CTR Interest, revised, ORNL-TM-3233, Oak Ridge National Laboratory, Oak Ridge TN, pp. 2-5, 1971.
- 27) Clark, D. A., et al., IEEE Transactions on Nuclear Science, Vol. NS-28, No. 2, p. 1339, April 1981.

- 28) Glasstone, G., and Loveberg, R. W., Controlled Thermonuclear Reactions, Robert E. Krieger Publishing Co., Huntington NY, 1960.
- 29) Tombrello, T. A., Recent $D^{11}(p,3He^4)$ Cross Section Requirements, W. K. Kellogg Radiation Laboratory, California Institute of Technology, Pasadena CA, 1977.
- 30) Wittenberg, L. J., Santarius, J. F., and Kulcinski, G. L., "Lunar Source of He^3 for Commercial Space Power," Fusion Technology, Vol. 10, p. 167, September 1986.
- 31) Norbeck, E., Cross Section Requirements and Needs, EPRI-536-SR, Proceedings of the Review Meeting on Advanced-Fuel Fusion, edited by K. Choi, Electric Power Research Institute, Palo Alto CA, p. 226-32, June 1977.
- 32) Feldbacher, R., The AEP Barnbook--DATLIB: Nuclear Reaction Cross Sections and Reactivity Parameter Library and Files, AEP Program, Institute for Theoretical Physics, Technical Univ. Graz, Petergasse, October 1987.
- 33) Hale, G. M., $D^{11}(p,3He^4)$ Reaction, evaluation of California Institute of Technology and Segel data, Los Alamos National Laboratory, Los Alamos NM, July 1979.
- 34) Feldbacher, R., Heindler, M., and Miley, G., Requirements for Charged Particle Light Isotopes Reaction Data for Advanced Fuel Cycles, Including Two Step Reaction Mechanism, Univ. of Illinois, Dept. of Nuclear Engineering, Urbana IL, February 1986.
- 35) Wittenberg, L. J., Santarius, J. F., and Kulcinski, G. L., Lunar Source of He^3 for Commercial Fusion Power, ANS Reprint, Fusion Technology, Vol. 10, pp. 167-78, September 1986.
- 36) Kulcinski, G. L., and Schmitt, H. W., The Moon: An Abundant Source of Clean and Safe Fusion Fuel for the 21st Century, UWFDN-730, Fusion Technology Institute, University of Wisconsin, Madison WI, August 1987.
- 37) Kulcinski, G. L., Santarius, J. F., and Wittenberg, L. J., Clean Thermonuclear Power from the Moon, UWFDN-709, Fusion Technology Institute, University of Wisconsin, Madison WI, September 1986.